


IN THE CLAIMS:

Amend the claims as follows.


Claims 1-28. (Canceled)

29. (New) Precursors of drugs with an anti-malarial action, characterized in that they are precursors of quaternary bis-ammonium salts and that they correspond to general formula (I)



in which

- A and A' are identical to or different from one another and represent either, an A₁ and A'₁ group respectively, of formula



where n is an integer from 2 to 4; R'₁ represents a hydrogen atom, a C1 to C5 alkyl radical, optionally substituted by an aryl radical, a hydroxy, an alkoxy, in which the alkyl radical comprises from 1 to 5 C, or aryloxy; and W represents a halogen atom chosen from chlorine, bromine or iodine, or a nucleofuge group, *selected from the group consisting of*


and
 C₆H₄-SO₃, mesityl CH₃-SO₃, CF₃-SO₃, NO₂-C₆H₄-SO₃ radical,


or an \underline{A}_2 group which represents a formyl -CHO radical, or an acetyl -CO-CH₃,

- \underline{B} and \underline{B}' are identical to or different from one another and represent

either the \underline{B}_1 and \underline{B}'_1 groups respectively, if \underline{A} and \underline{A}' represent \underline{A}_1 and \underline{A}'_1 respectively, \underline{B}_1 and \underline{B}'_1 representing an \underline{R}_1 group which has the same definition as \underline{R}'_1 above, but cannot be a hydrogen atom,

or the \underline{B}_2 and \underline{B}'_2 groups respectively, if \underline{A} and \underline{A}' represent \underline{A}_2 , \underline{B}_2 or \underline{B}'_2 being the \underline{R}_1 group as defined above, or a group of formula




in which -Ra represents an RS- or RCO- group, where \underline{R} is a linear, ^{branched} branched or  cyclic C1 to C6 alkyl radical, optionally substituted by one or more hydroxy, alkoxy or aryloxy group, or an amino group and/or a -COOH or COOM group, where \underline{M} is a C1 to C3 alkyl; a phenyl or benzyl radical, in which the phenyl radical is optionally substituted by at least one C1 to C5 alkyl or alkoxy radical, these being optionally substituted by an amino group, or by a nitrogenous or oxygenous heterocycle, a -COOH or -COOM group; or a saturated -CH₂-heterocycle group, with 5 or 6 elements, nitrogenous and/or oxygenous; \underline{R}_2 represents a hydrogen atom, a C1 to C5 alkyl radical, or a -CH₂-COO-alkyl (C1 to C5) group; and \underline{R}_3 represents a hydrogen atom, a C1 to C5 alkyl or alkenyl radical, optionally substituted by -OH, a phosphate group, an alkoxy radical, in which the alkyl radical is C1 to C3, or an aryloxy radical; or an alkyl (or aryl), carbonyloxy


group; or also \underline{R}_2 and \underline{R}_3 together form a ring with 5 or 6 carbon atoms; R and \underline{R}_3 can be linked to form a cycle of 5 to 7 atoms (carbon, oxygen, sulphur)

- $\underline{\alpha}$ represents


either a single bond, when \underline{A} and \underline{A}' represent \underline{A}_1 and \underline{A}'_1 ; or when \underline{A} and \underline{A}' represent \underline{A}_2 , i.e. a -CHO or -COCH₃ group, and \underline{B}_2 and \underline{B}'_2 represent



or, when \underline{A} and \underline{A}' represent \underline{A}_2 and \underline{B}_2 and \underline{B}'_2 represent \underline{R}_1 , a group of formula



or a group of formula




in which (a) represents a bond towards \underline{Z} and (b) a bond towards the nitrogen atom.

- \underline{Z} represents a C6 to C21 alkylene radical, optionally with insertion of one or more multiple bonds, and/or one or more O and/or S heteroatoms, and/or one or more aromatic rings, and the pharmaceutically acceptable salts of these compounds,

provided that R'_1 does not represent H or a C1 or C2 alkyl radical, when $n = 3$ or 4 , R_1 represents a C1 to C4 alkyl radical and Z represents a C6 to C10 alkylene radical.

²
~~30.~~ (New) Precursors according to claim ¹~~29~~, characterized in that it relates to haloalkylamines, corresponding to general formula (II)



in which R_1 , R'_1 , W , n and Z are as defined in claim 29.

³
~~31.~~ (New) Precursors according to claim ¹~~29~~, wherein Z represents a C13 to C21 alkylene radical.


⁴
~~32.~~ (New) Precursors according to claim ³~~31~~, characterized in that Z represents a $-(CH_2)_{16}$ - group.

⁵
~~33.~~ (New) Precursors according to claim ²~~30~~ or ³~~31~~, characterized in that R_1 is a methyl radical.


⁶
~~34.~~ (New) Precursors according to claim ²~~30~~, characterized in that R_1 is a methyl radical and R'_1 is either a hydrogen atom, or a methyl radical, and W is a chlorine atom.

3
7 35. (New) Precursors according to claim 31, characterized in that they are chosen from N, N'-dimethyl-N,N'-(5-chloropentyl)-1,16-hexadecanediamine hydrochloride, or N, N'-dimethyl-N,N'-(4-chloropentyl)-1,16-hexadecanediamine hydrochloride.


8 36. (New) Precursors according to claim 29, characterized in that it concerns precursors of thiazolium corresponding to general formula (III).



or to general formula (IV)



or to general formula (V)



in which $\underline{R_a}$, $\underline{R_1}$, $\underline{R_2}$, and \underline{Z} are as defined in claim 29.

9
~~37~~. (New) Precursors according to claim ~~36~~⁸, characterized in that they correspond to formula III in which $\underline{R_a}$ represents an RCO- radical.

10
~~38~~. (New) Precursors according to claim ~~37~~⁹, characterized in that they are chosen from N,N'-diformyl-N,N'-di[1-methyl-2-S-thiobenzoyl-4-methoxybut-1-enyl]-1,12-diaminododecane, N,N'-diformyl-N,N'-di[1-methyl-2-S-(p-diethylaminomethylphenylcarboxy)thio-4-methoxybut-1-enyl]-1,12-diaminododecane, N,N'-diformyl-N,N'-di[1-methyl-2-S-(p-morpholino-methylphenylcarboxy)-thio-4-methoxybut-1-enyl]-1,12-diaminododecane, N,N'-diformyl-N,N'-di[1-methyl-2-S-thiobenzoyl-4-methoxybut-1-enyl]-1,16-diaminohexadecane, and N,N'-diformyl-N,N'-di[1(2-oxo-4,5-dihydro-1,3-oxathian-4-ylidene)ethyl]1,12-diaminododecane.


11
~~39~~. (New) Precursors according to claim ~~37~~⁹, characterized in that $\underline{R_a}$ represents RS-.

²40. (New) Precursors according to claim ¹¹39, characterized in that they are chosen from N,N'-diformyl-N,N'-di[1-methyl-2-tetrahydrofurfuryl-methyldithio-4-hydroxybut-1-enyl]-1,12-diaminododecane, N,N'-diformyl-N,N'-di[1-methyl-2-propyldithio-4-hydroxybut-1-enyl]-1,12-diaminododecane, N,N'-diformyl-N,N'-di[1-methyl-2-benzyl-dithio-4-hydroxybut-1-enyl]-1,12-diaminododecane, N,N'-diformyl-N,N'-di[1-methyl-2-(2-hydroxyethyl)-dithio-4-hydroxybut-1-enyl]-1,12-diaminododecane, N,N'-diformyl-N,N'-di[1-methyl-2-propyldithio-4-methoxybut-1-enyl]-1,12-diaminododecane, and N,N'-diformyl-N,N'-di[1-methyl-2-propyldithio-ethenyl]-1,12-diaminododecane.

³41. (New) Precursors according to claim ⁸36, characterized in that they correspond to formula IV and are chosen from 2,17-(N,N'-diformyl-N,N'-dimethyl)diamino-3,16-S-thio-p-methoxybenzoyl-6,13-dioxaoctadeca-2,16-diene, 2,17-(N,N'-diformyl-N,N'-dibenzyl)diamino-3,16-S-thio-p-methoxybenzoyl-6,13-dioxaoctadeca-2,16-diene, ethyl 3,18 (N,N'-diformyl-N,N'-dimethyldiamino-4,17-S-thiobenzoyl-eicosa-3,17-dienedioate (TE12), ethyl 3,18-(N,N'-diformyl-N,N'-dibenzyl)diamino-4,17-S-thiobenzoyl-eicosa-3,17-dienedioate.


¹⁴42. (New) Precursors according to claim ⁸36, characterized in that they correspond to formula (V) and are chosen from 2,15-(N,N'-diformyl-N,N'-dimethyl)diamino-1,16-S-thiobenzoyl-hexadeca-1,15-diene. 2,15-(N,N'-diformyl-N,N'-dibenzyl)diamino-1,16-S-thio-benzoyl-hexadeca-1,15-diene.

43. (New) The cyclized derivatives corresponding to the precursors of thiazolium of claim 29, said derivatives having general formula (VI)




in which

R_b represents R₁ or T, T representing the group of formula




provided that Z does not represent a C6 to C8 alkylene radical, when R_c, R_d, R₂ and R₃ represent a methyl radical.

R_d represents R₂ or P, P representing the group of formula



R_c represents R₃ or U, U representing the group of formula



R₁, R₂, R₃ and Z being as defined in claim 29,

it being understood that $\underline{R}_b = \underline{T}$, if $\underline{R}_c = \underline{R}_3$ and $\underline{R}_d = \underline{R}_2$; $\underline{R}_d = \underline{P}$, if $\underline{R}_c = \underline{R}_3$ and $\underline{R}_b = \underline{R}_1$; and $\underline{R}_c = \underline{U}$, if $\underline{R}_b = \underline{R}_1$ and $\underline{R}_d = \underline{R}_2$.


¹⁶
~~44~~. (New) Process for obtaining precursors of thiazolium of general formula (III) to (IV) according to claim ⁷~~35~~, characterized in that it comprises the reaction in basic medium of a thiazole derivative of formula (VI).

¹⁷
~~45~~. (New) Process according to claim ¹⁶~~44~~, characterized in that in order to obtain the compounds in which $\underline{R}_a = \text{RCO}-$, a derivative of thiazolium of formula (VI) is reacted with an RCOR' derivative, where \underline{R} is as defined in claim 1 and \underline{R}' is a halogen atom, and in order to obtain the compounds in which $\underline{R}_a = \text{RS}-$, said thiazolium derivatives of formula (VI) are reacted with a thiosulphate derivative $\text{RS}_2\text{O}_3\text{Na}$.

¹⁸
~~46~~. (New) Process according to claim ¹⁵~~43~~ or ¹⁶~~44~~, characterized in that


- in order to obtain the compounds of formula (III) a thiazole derivative suitably substituted with an alkyl dihalide is reacted, under reflux in an organic solvent, the opening of the thiazolium ring then takes place in basic medium, and by the action either of R-COCl , or of $\text{RS}_2\text{O}_3\text{Na}$,

- in order to obtain the compounds of formula IV, which comprise an oxygen in the \underline{Z} chain, a thiazole derivative of general formula (VII)



is reacted with an alkane dihalide, in basic medium, then the addition of R_1X , the reaction medium being advantageously taken to reflux in an organic solvent, in particular alcoholic such as ethanol, for a duration sufficient to obtain the quaternization of the nitrogen atom of the thiazole by fixing R_1 , the opening of the thiazolium ring then being obtained in basic medium, then by the action either of $R-COCl$, or of RS_2O_3Na ,


- in order to obtain the compounds of formula (IV) not comprising oxygen in the Z chain, a compound of structure



is firstly synthesized by reacting an alkyl acetoacetate with NaH , followed by alkylation, then the addition of a dihalogenoalkane, the compound obtained then being dibrominated, then thioformamide is added and, after reflux for several days, R_1X , which leads, after renewed reflux for several days, to a thiazolium, the opening of which is then carried out in basic medium, then the action of $R-COCl$ or of $R-S_2O_3Na$,


- in order to obtain the compounds of formula (V) not comprising oxygen in the Z chain, a $Z(CO-CH_2 X)_2$ compound is reacted with $CH(=S)NH_2$, then R_1X is added, the opening of the thiazolium ring then being carried out in basic medium, then by adding $R-COCl$ or $R-S_2O_3Na$.

47. (New) Process for obtaining haloalkylamines according to claim 29, characterized in that it comprizes the alkylation of an amino alcohol of formula




by an alkyl α,ω -dihalide X-Z-X, which leads to a bis-aminoalcohol treated with a compound capable of releasing the W group.

48. (New) Pharmaceutical compositions, characterized in that they contain an effective quantity of at least one precursor as defined in claim 29, or at least one cyclized derivative of general formula (VI):




in which


R_b represents R_1 or I , I representing the group of formula:



R_d represents R_2 or P , P representing the group of formula



R_c represents R₃ or U, U representing the group of formula




R₁, R₂, R₃ and Z being as defined in claim 29,

it being understood that R_b = T, if R_c = R₃ and R_d = R₂; R_d = P, if R_c = R₃ and R_b = R₁; and R_c = U, if R_b = R₁ and R_d = R₂.

in combination with a pharmaceutically inert vehicle.


2
 49. (New) A method of treating at least one of malaria and babesiosis comprising administering to an animal in need of said treatment a quaternary bis-ammonium salt of formula I



in which

- A and A' are identical to or different from one another and represent

either, an A₁ and A'₁ group respectively, of formula




where n is an integer from 2 to 4; R₁' represents a hydrogen atom, a C1 to C5 alkyl radical, optionally substituted by an aryl radical, a hydroxy, an alkoxy, in which the alkyl radical comprises from 1 to 5 C, or aryloxy; and W represents a halogen atom chosen from chlorine, bromine or iodine, or a nucleofuge group, such as the tosyl CH₃-C₆H₄-SO₃, mesityl CH₃-SO₃, CF₃-SO₃, NO₂-C₆H₄-SO₃ radical,

or an A₂ group which represents a formyl -CHO radical, or an acetyl-CO CH₃ radical,

- B and B' are identical to or different from one another and represent

either the B₁ and B₁' groups respectively, if A and A' represent A₁ and A₁' respectively, B₁ and B₁' representing an R₁ group which has the same definition as R₁' above, but cannot be a hydrogen atom,

or the B₂ and B₂' groups respectively, if A and A' represent A₂, B₂ or B₂' being the R₁ group as defined above, or a group of formula




in which -R_a represents an RS- or RCO- group, where R is a linear, branched or cyclic C1 to C6 alkyl radical, optionally substituted by one or more hydroxy, alkoxy or


aryloxy group, or an amino group and/or a -COOH or COOM group, where M is a C1 to C3 alkyl; a phenyl or benzyl radical, in which the phenyl radical is optionally substituted by at least one C1 to C5 alkyl or alkoxy radical, these being optionally substituted by an amino group, or by a nitrogenous or oxygenous heterocycle, a -COOH or -COOM group; or a saturated -CH₂-heterocycle group, with 5 or 6 elements, nitrogenous and/or oxygenous; R₂ represents a hydrogen atom, a C1 to C5 alkyl radical, or a -CH₂-COO-alkyl (C1 to C5) group; and R₃ represents a hydrogen atom, a C1 to C5 alkyl or alkenyl radical, optionally substituted by -OH, a phosphate group, an alkoxy radical, in which the alkyl radical is C1 to C3, or an aryloxy radical; or an alkyl (or aryl), carbonyloxy group; or also R₂ and R₃ together form a ring with 5 or 6 carbon atoms; and R and R₃ can be linked to form a cycle of 5 to 7 atoms (carbon, oxygen, sulphur)

- α represents


either a single bond, when A and A' represent A₁ and A₁': or when A and A' represent A₂, i.e. a -CHO or -COCH₃ group, and B₂ and B₂' represent



or, when A and A' represent a -CHO group and B₂ and B₂' represent R₁, a group of formula



or a group of formula



in which (a) represents a bond towards \underline{Z} and (b) a bond towards the nitrogen atom.

- \underline{Z} represents a C6 to C21 alkylene radical, optionally with insertion of one or more multiple bonds, and/or one or more O and/or S heteroatoms, and/or one or more aromatic rings, and the pharmaceutically acceptable salts of these compounds.

²²
50. (New) Pharmaceutical compositions according to claim ²⁰~~48~~, in a form which may be administered by at least one of the oral route, injectable route, or rectal route.

²³
51. (New) A precursor according to claim ¹~~29~~ wherein said aryl radical is a phenyl radical.

²⁴
52. (New) A method according to claim ²¹~~49~~ wherein said aryl radical is a phenyl radical.

²⁵
53. (New) A precursor according to claim ¹~~29~~ wherein said aryloxy is a phenoxy.

²⁶
54. (New) A method according to claim ²¹~~49~~ wherein said aryloxy is a phenoxy.

²⁷
55. (New) A method according to claim ²¹~~49~~ wherein said animal is a man.